

**UNCLASSIFIED**

---

**AD 296 898**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

296 898

# OPTIMALITY PROPERTIES

OF A SPECIAL ASSIGNMENT PROBLEM

S. C. J.  
Opera  
Universit

by

H. Parikh and Roger Wets  
Operations Research Center  
University of California, Berkeley

3 December 1962

Research Report

This research was primarily  
Institute of the University of  
from the Office of Naval Re  
Science Foundation Research  
ornia, Berkeley. Reproduc  
pose of the United States Go

ly supported by the Western Management Scien  
of California, Los Angeles, with partial supp  
Research, Contract Nonr-222(83) and the Natio  
Grant G-21034, with the University of Cal  
tion in whole or in part is permitted for any p  
government.

## ABSTRACT

In this paper, it is shown that if the cost matrix of an assignment problem has the following property  $c_{ij} = |j - i|$  then any basic feasible solution is optimal if and only if its unit components belong to two well defined symmetric regions. The matrix with above mentioned property is called the "REORDERING MATRIX," because it arose for the first time in the reordering of nodes of a critical path and other acyclic network problems.

## OPTIMALITY PROPERTIES OF A SPECIAL ASSIGNMENT PROBLEM

### I. Introduction

Assume we want to order the nodes of a network in such a way that for every arc  $(i, j)$ , which belongs to the topology of the network,  $i < j$ .<sup>\*</sup> In case of a large network, this may not be possible to do manually, as it generates a large permutation problem. An algorithm to reorder the nodes of the network is given in [1], where the number of steps involved is related to the magnitude of divergence of the node order of a network, where divergence is defined to be

$$S = \sum_{i=1}^m |i - a_i|$$

where  $a_i$  is the initial order of the node having final order  $i$ .

In order to fix an upper bound to the divergence of the network, we have to determine the maximum value of  $S$  over all possible  $m$ -tuples  $a$ .

We will now show that this problem is equivalent to the classical assignment problem.

---

<sup>\*</sup>This problem is important because it reduces significantly the work involved in logical search for a critical path problem. It is impossible to perform this node ordering when loops are present in the network.

## II. Mathematical Formulation

Let  $\mathcal{A} = \left\{ a \mid a \text{ is an } m\text{-tuple}^* \text{ chosen without repetition from the set } \{1, 2, \dots, m\} \text{ e.g., } a = (2, m, 10, \dots, 1, 5) \right\}$ .

Note that  $\mathcal{A}$  has  $m!$  nonidentical components.

Then, the problem becomes

$$(1) \quad \max_{a \in \mathcal{A}} S = \sum_{i=1}^m |i - a_i|,$$

where the  $m$ -tuples " $a$ " are chosen without repetition from the set of  $m$  first integers, to each integer  $k$  corresponding one and only one  $a_i$  for each  $m$ -tuple " $a$ ". It is now easy to see that all the components of each possible summation  $\sum_{i=1}^m |i - a_i|$  are of the form  $|i - j|$  where  $i = 1, \dots, m$ ;  $j = 1, \dots, m$ . We can list all these possible values under matrix form where each entry is equal to the absolute difference between its row number and column number.

Define  $c_{ij} = |i - j|$  and  $C = \{c_{ij}\}$ . Any possible  $S$  can be found by summing up  $m$  entries of the matrix selected by picking one and only one element from each row and each column. For example

i	$a_i = j$			
1	$c_{11}$	$c_{12}$	$c_{13}$	$c_{14}$
2	$c_{21}$	$c_{22}$	$c_{23}$	$c_{24}$
3	$c_{31}$	$c_{32}$	$c_{33}$	$c_{34}$
4	$c_{41}$	$c_{42}$	$c_{43}$	$c_{44}$

\*  $m$ -tuple " $a$ " has  $m$  elements, where  $a_i$  ( $i = 1, \dots, m$ ) is the  $i^{\text{th}}$  element.

A possible  $S = c_{12} + c_{23} + c_{31} + c_{44}$  corresponding to the 4-tuple "a" = (2, 3, 1, 4) .

Determining the  $\max_{a \in A} S$  is equivalent to finding some feasible combination of the entries of the C matrix (i.e., one and only one entry in each row and each column) such that their sum is maximum. Thus, the problem reduces to

$$\text{maximize } S = \sum_{ij} c_{ij} x_{ij}$$

Subject to:

$$\sum_j x_{ij} = 1 \quad i = 1, \dots, m$$

$$(2) \quad \sum_i x_{ij} = 1 \quad j = 1, \dots, m$$

$$x_{ij} \geq 0 \quad x_{ij} \text{ are integers ,}$$

which is the classical assignment problem formulation.

Our aim is to prove, in this particular set-up, that this optimum for S can be achieved by selecting any feasible solution such that all its components belong to two symmetric regions of the matrix and that no optimal solution can be found if one or more components do not belong to these two regions.

### III. The Optimal Region

In the reordering matrix let us define the two following sets of entries.

$$O_1^* = \{c_{ij} \mid i \leq \frac{m+1}{2} , j \geq \frac{m+1}{2} \}$$

$$O_2^* = \{c_{ij} \mid i \geq \frac{m+1}{2} , j \leq \frac{m+1}{2} \} .$$

Let  $O^* = O_1^* \cup O_2^*$  then,

$$O_1^* \cap O_2^* = \begin{cases} \emptyset & \text{if } m \text{ is even} \\ c_{m+1/2, m+1/2} & \text{if } m \text{ is odd} \end{cases} .$$

These two sets of entries are symmetric with respect to the principal diagonal




Figure 1 – Optimal Region.

$$\text{Let } O_1 = \{x_{ij} \mid x_{ij} \in O_1 \text{ iff } c_{ij} \in O_1^*\}$$

$$O_2 = \{x_{ij} \mid x_{ij} \in O_2 \text{ iff } c_{ij} \in O_2^*\} .$$

Then,  $O = O_1 \cup O_2$  and it is called the optimal region.



NOTE: If we were minimizing  $\sum c_{ij}x_{ij}$  then the optimal solution would be unique and would be  $x = (x_{11}, x_{22}, x_{33}, \dots, x_{mm})$  which is easy to see because all the elements of the principal diagonal are zero and the other elements are strictly positive.

#### IV. Properties of the Reordering Matrix

1. For all  $c_{rs} \notin O^*$

either all  $c_{rk} \in O^*$  are greater than or equal to  $c_{rs}$   
or all  $c_{ks} \in O^*$  are greater than or equal to  $c_{rs}$ .

2. For any submatrix of  $C$  of the form

$$C_s = \begin{bmatrix} c_{ij} & c_{i,j+1} \\ c_{i+1,j} & c_{i+1,j+1} \end{bmatrix}$$

such that  $i \neq j$

then  $c_{ij} + c_{i+1,j+1} = c_{i+1,j} + c_{i,j+1}$ . In fact this property is true for any submatrix chosen so that all its elements

$$C_s = \begin{bmatrix} c_{ij} & c_{i,j+s} \\ c_{i+r,j} & c_{i+r,j+s} \end{bmatrix}$$

are all below the principal diagonal or above the principal diagonal.

3. For any submatrix of  $C$  of the form

$$C_r = \begin{bmatrix} c_{ij} & c_{i,j+s} \\ c_{i+r,j} & c_{i+r,j+s} \end{bmatrix}$$

such that  $c_{i+r,j}$  is an entry below the principal diagonal of  $C$  and  $c_{i,j+s}$  is an entry above the principal diagonal. Then

$$c_{ij} + c_{i+r,j+s} < c_{i+r,j} + c_{i,j+s}$$

PROOF: When  $c_{i+r,j}$  is below the principal diagonal of  $C$ , it implies that

$$(3) \quad i + r > j \Rightarrow i - j + r > 0 \Rightarrow r > j - i .$$

When  $c_{i,j+s}$  is above the principal diagonal of  $C$ , it implies that

$$(4) \quad j + s > i \Rightarrow i - j - s < 0 \Rightarrow s > i - j .$$

$c_{ij} + c_{i+r,j+s} < c_{i+r,j} + c_{i,j+s}$  can be written as follows:

$$|i - j| + |i - j + r - s| < |i - j + r| + |i - j - s| .$$

From (3) and (4), we get

$$(5) \quad \begin{aligned} |i - j| + |i - j + r - s| &< (i - j + r) - (i - j - s) , \\ |i - j| + |i - j + r - s| &< r + s . \end{aligned}$$

Here we can distinguish various cases. We will prove it for one case.

Assume  $i > j$  and  $r > s$ ; (5) then becomes

$$i - j + i - j + r - s < r + s$$

then

$$2i - 2j < 2s$$

or

$$i - j < s$$

which is relation (4).

It is possible to prove the optimality of any feasible solution in region  $O$  by using the theory of linear programming, or more specifically the assignment problem algorithm. [2] Let the simplex multipliers be

$$\begin{aligned} \pi &= -i \quad \text{for the constraints} \quad \sum_j x_{ij} = 1 && \text{for } i < \frac{m+1}{2} \\ &= +i \quad \text{for the constraints} \quad \sum_j x_{ij} = 1 && \text{for } i \geq \frac{m+1}{2} \\ \pi &= -j \quad \text{for the constraints} \quad \sum_i x_{ij} = 1 && \text{for } j < \frac{m+1}{2} \\ &= +j \quad \text{for the constraints} \quad \sum_i x_{ij} = 1 && \text{for } j \geq \frac{m+1}{2} . \end{aligned}$$

If we price out, it is possible to see that the  $\bar{c}_{ij} \in O^*$  are equal to zero and the other  $\bar{c}_{ij} \notin O^*$  are <sup>negative</sup> ~~positive~~. We are going to give an alternative proof which will use the properties of the reordering matrix and the fact that the solution has to be feasible.

#### V. Optimality Theorem

**THEOREM:** A feasible solution  $x^0$  is optimal if and only if all its components lie in  $O$ .

**PROOF:** Suppose  $\hat{x}$  is a feasible solution but has at least one component which does not lie in  $O$ , we will then show that we can improve  $S$ .

Let us assume that  $\hat{x}_{ij}$  is such that  $i < \frac{m+1}{2}$ ,  $j < \frac{m+1}{2}$  and  $j \geq i$ , i.e., in Figure 2,  $\hat{x}_{ij}$  lies in  $U$ . In order to be feasible  $x$  has at least a component, say  $\hat{x}_{kl} \in V$ , because it is impossible to "cover"




Figure 2

the  $\left\lfloor \frac{m+1}{2} \right\rfloor$  \* last rows with "selections" only done in  $V$ , as  $V$  has less than  $\left\lfloor \frac{m+1}{2} \right\rfloor$  columns (by assumption on  $\hat{x}_{ij}$ ). We can then find a new feasible solution by replacing  $\hat{x}_{ij}$  and  $\hat{x}_{kl}$  by  $x_{il}$  and  $x_{kj}$  and by property 3,  $S$  will be improved.

We have proved that for any feasible solution which has one or more components outside  $O$ , it can be improved. So, we can produce an iterative procedure which will increase  $S$ , as long as  $x$  has a component which does not lie in  $O$ .

If  $x^0 \in O$ , it is not possible to improve the value of  $S$  because the only acceptable substitutions are of the form

$$\begin{array}{l} x_{ij}^0 \quad \text{and} \quad x_{i+r,j+s}^0 \\ \text{by} \quad x_{i+r,j}^0 \quad \text{and} \quad x_{i,j+s}^0 \quad (\in O) \end{array}$$

\*  $[\alpha]$  = greatest integer contained in  $\alpha$ .

or repeated substitutions of that form. But by property 2 we know that the value of  $S$  will not change. If one of the components of our substitutions does not lie in  $O$  then we have proved that we can improve the value of  $S$ .

**COROLLARY:** All the feasible solutions in region  $O$  are optimal.

**PROOF:** By property 2 of the reordering matrix, from an optimal solution  $x^O$ , we can find new optimal solutions  $\bar{x}$  by repeated substitutions of the form:

$$(6) \quad x_{ij}^O = x_{i+r,j+s}^O = 1, \quad x_{i+r,j}^O = x_{i,j+s}^O = 0$$

by

$$\bar{x}_{ij} = \bar{x}_{i+r,j+s} = 0, \quad \bar{x}_{i+r,j} = x_{i,j+s} = 1,$$

as long as  $x_{i+r,j}$  and  $x_{i,j+s}$  belong to  $O$ .

**NOTE:** By repeated substitution of the form (6) it is possible to reproduce all  $m$ -tuples  $a \in \mathcal{A}$  in matrix  $C$ .

## REFERENCES

- [ 1 ]. S. Parikh, "Project Planning by Decomposition, " Research Report ,  
Operations Research Center, University of California, Berkeley,  
(To be published.)
- [ 2 ]. L.R. Ford and D.R. Fulkerson, "Flows in Networks, " Princeton  
University Press, Princeton, New Jersey, 1962.

# BASIC DISTRIBUTION LIST FOR UNCLASSIFIED TECHNICAL REPORTS

Head, Logistics and Mathematical  
Statistics Branch  
Office of Naval Research  
Washington 25, D. C.

C. O., ONR Branch Office  
Navy No. 100, Box 39, F. P. O.  
New York City, New York

ASTIA Document Service Center  
Arlington Hall Station  
Arlington 12, Virginia

Institute for Defense Analyses  
Communications Research Div.  
von Neumann Hall  
Princeton, New Jersey

Technical Information Officer  
Naval Research Laboratory  
Washington 25, D. C.

C. O., ONR Branch Office  
346 Broadway, New York 13, NY  
Attn: J. Laderman

C. O., ONR Branch Office  
1030 East Green Street  
Pasadena 1, California  
Attn: Dr. A. R. Laufer

Bureau of Supplies and Accounts  
Code OW, Dept. of the Navy  
Washington 25, D. C.

Professor Russell Ackoff  
Operations Research Group  
Case Institute of Technology  
Cleveland 6, Ohio

Professor Kenneth J. Arrow  
Serra House, Stanford University  
Stanford, California

Professor G. L. Bach  
Carnegie Institute of Technology  
Planning and Control of Industrial  
Operations, Schenley Park  
Pittsburgh 13, Pennsylvania

Professor A. Charnes  
The Technological Institute  
Northwestern University  
Evanston, Illinois

Professor L. W. Cohen  
Math. Dept., University of Maryland  
College Park, Maryland

Professor Donald Eckman  
Director, Systems Research Center  
Case Institute of Technology  
Cleveland, Ohio

Professor Lawrence E. Fouraker  
Department of Economics  
The Pennsylvania State University  
State College, Pennsylvania

Professor David Gale  
Dept. of Math., Brown University  
Providence 12, Rhode Island

Dr. Murray Geisler  
The RAND Corporation  
1700 Main Street  
Santa Monica, California

Professor L. Hurwicz  
School of Business Administration  
University of Minnesota  
Minneapolis 14, Minnesota

Professor James R. Jackson  
Management Sciences Research  
Project, Univ. of California  
Los Angeles 24, California

Professor Samuel Karlin  
Math. Dept., Stanford University  
Stanford, California

Professor C. E. Lemke  
Dept. of Mathematics  
Rensselaer Polytechnic Institute  
Troy, New York

Professor W. H. Marlow  
Logistics Research Project  
The George Washington University  
707 - 22nd Street, N. W.  
Washington 7, D. C.

Professor Oskar Morgenstern  
Economics Research Project  
Princeton University  
92 A Nassau Street  
Princeton, New Jersey

**BASIC DISTRIBUTION LIST  
FOR UNCLASSIFIED TECHNICAL REPORTS**

**Professor R. Radner  
Department of Economics  
University of California  
Berkeley, California**

**Professor Stanley Reiter  
Department of Economics  
Purdue University  
Lafayette, Indiana**

**Professor Murray Rosenblatt  
Department of Mathematics  
Brown University  
Providence 12, Rhode Island**

**Mr. J. R. Simpson  
Bureau of Supplies and Accounts  
Navy Department (Code W31)  
Washington 25, D.C.**

**Professor A. W. Tucker  
Department of Mathematics  
Princeton University  
Princeton, New Jersey**

**Professor J. Wolfowitz  
Department of Mathematics  
Lincoln Hall, Cornell University  
Ithaca 1, New York**